Check List the journal of biodiversity data

NOTES ON GEOGRAPHIC DISTRIBUTION

Check List 17 (2): 339–345
https://doi.org/10.15560/17.2.339

PENSOFT

Altitudinal range extension of Long-tailed Singing Mouse, *Scotinomys xerampelinus* (Bangs, 1902) (Rodentia, Cricetidae), in Chirripó National Park, Costa Rica

José Manuel Mora[1,2*], Jairo García Céspedes[3], Lucía Isabel López[4], Gerardo Chaves[5]

1 Carrera de Gestión Ecoturística, Sede Central y Unidad de Ciencias Básicas, Sede Atenas, Universidad Técnica Nacional, Alajuela, Costa Rica • josemora07@gmail.com https://orcid.org/0000-0002-1200-1495

2 Maestría en Desarrollo Sostenible, Sede de Occidente, Universidad de Costa Rica, San Ramón, Costa Rica

3 Laboratorio de Geoquímica, Escuela Centroamericana de Geología, Universidad de Costa Rica, San José, Costa Rica • jairo.garcia@ucr.ac.cr

4 Unidad de Ciencias Básicas y Carrera de Tecnología de Alimentos, Sede Atenas, Universidad Técnica Nacional, Atenas, Costa Rica https://orcid.org/0000-0002-0120-7981

5 Museo de Zoología, Universidad de Costa Rica, San Pedro de Montes de Oca, Costa Rica • cachi13@gmail.com https://orcid.org/0000-0002-4301-6569

* Corresponding author

Abstract

Scotinomys xerampelinus has a restricted distribution in the Cordilleras Central and Talamanca of Costa Rica and western Panama, at an elevational range between 2100 and 3400 m. We report individuals observed at Cerro Chirripó, Costa Rica at 3820 m, which extends upwards the known elevational range by 420 m. The altitudinal range extension may indicate either incomplete surveys in the study area or an upslope shift due to increasing temperatures from climate change, a phenomenon that has forced several Costa Rican vertebrate species to transition to higher elevations.

Keywords

Cloud forest, montane rain forest, Neotropics, Páramo, rodents, small mammals

Academic editor: Héctor Ramírez-Chaves | Received 5 September 2020 | Accepted 25 January 2021 | Published 2 March 2021

Citation: Mora JM, Céspedes JG, López LI, Chaves G (2021) Altitudinal range extension of Long-tailed Singing Mouse, *Scotinomys xerampelinus* (Bangs, 1902) (Rodentia, Cricetidae), in Chirripó National Park, Costa Rica. Check List 17 (2): 339–345. https://doi.org/10.15560/17.2.339

Introduction

Biodiversity is distributed globally based on systematic patterns of spatial variation (Pianka 1989). In general, species richness tends to be correlated with latitude and elevation gradients, with the latter occurring across smaller spatial scales (Hillebrand 2004; McCain 2007; but see Wen et al. 2021). Several hypotheses have tried to explain these associations based on, for example, the effects of availability of energy or temperature on population dynamics coupled with the time available for the process of diversification to occur (Allen et al. 2002; Dyer et al. 2020). The complex interrelatedness of climate on trends in biodiversity has been recognized as important in altitudinal distributions (Brown 2001). Considering the changing global climate, it is urgent to understand how temperature changes affect all types of biological systems, from individuals to whole ecosystems

©The authors. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

(Kontopoulos et al. 2020). At the same time, distributional changes of animal and plant species is a matter of great concern for biodiversity conservation (Mora and López 2014).

Costa Rica is a small country (51100 km^2) and covers only 0.034% of the land surface of the Earth. However, with over 250 mammal species, this country contains approximately 5% of the world's known mammal species (SINAC 2014). In the last two decades, knowledge of Costa Rican mammals has notably increased (Rodríguez-Herrera et al. 2014; Villalobos-Chaves et al. 2016; Mora et al. 2020), although much remains unknown. One of the least known groups in the country is rodents, and frequently even distributional limits of species are unknown (Rodríguez-Herrera et al. 2014). Only a few publications constitute the basis for knowledge of rodents and other mammals in Costa Rica. The pioneer publications by Frantzius (1869), Alfaro (1897), and Goodwin (1946) are among the most notable (Rodríguez-Herrera et al. 2005). At the beginning of the 1960s, the distributions of numerous species remained poorly known, notwithstanding the expansive application to Central America by Ryan (1963) of Dice's concept of biotic provinces (McPherson 1985). McPherson (1971a, 1971b, 1985) contributed greatly to the knowledge of the distribution ranges and ecological relationships of rodents in Costa Rica. The list of mammals of Costa Rica including new records and new information on distributional ranges of some species was later updated by subsequent authors (Wilson 1983; Mora 2000; Wilson et al. 2002; Rodriguez-Herrera et al. 2002, 2014). The most complete list of rodents of Costa Rica was published several years later (Villalobos-Chaves et al. 2016), and one additional rodent species, *Ichthyomys tweedii* Anthony, 1921, has been recently added (Ramírez-Fernández et al. 2020). However, all these publications include limited information and are concerned merely with presence/absence data; knowledge on the distributions of most rodents in Costa Rica is incomplete, and numerous gaps remain in the spatial distribution of most of small rodent species in the country.

Long-tailed Singing Mouse, *Scotinomys xerampelinus* (Bangs, 1902), is a cricetid rodent (Musser and Carleton 2005) which inhabits high elevations in the Cordilleras Central and Talamanca of southern Costa Rica and western Panamá between 2100 and 3400 m elevation (Naylor and Roach 2016). It is restricted to forests, meadows, and grasslands in two life zones in the high mountains: the montane rainforest and the páramo (Hooper and Carleton 1976). The páramo is a Neotropical high-mountain ecological association above the timberline consisting of ground cover by shrubs and grasses (Kappelle and Horn 2016). These two life zones are perennially wind-swept, cool, and draped in clouds or rain (Hooper and Carleton 1976). *Scotinomys xerampelinus* typically prefers areas having dense vegetative cover with mossy trunks, rocks, and dense fern cover (Bradley 2017). It is terrestrial and travels in runways under logs and among rocks (Hooper 1972). It is entirely diurnal and appears to eat only insects (Bradley 2017), mainly larval beetles which are probably located by smell (Reid 2009).

Males of *S. xerampelinus* are recognized by their vocalizations (Mora 2000) which they use to attract mates and to warn off other males from their territories (Pasch et al. 2013). Their songs also serve to repel a smaller, competing species, Alston's Singing Mouse, *Scotinomys teguina* (Alston, 1877) (Pasch et al. 2013). Long-tailed Singing Mouse is behaviorally dominant and excludes and abruptly replaces *S. teguina* at higher, cooler elevations (Hooper 1972; Hooper and Carleton 1976; Pasch et al. 2013). Here we report a new altitudinal record of Long-tailed Singing Mouse from the top of Chirripó mountain, Chirripó National Park, Costa Rica.

Methods

An unexpected encounter while visiting Cerro Chirripó, Parque Nacional Chirripó (PNC, Chirripó National Park in English) allowed us to photograph, measure, and release an individual *Scotinomys xerampelinus*. The PNC is located in the central section of the Cordillera de Talamanca (Veas-Ayala et al. 2018), a mountainous system with a southeast–northwest orientation that has the highest elevations in Costa Rica and separates the Caribbean plains from the lowlands of the Pacific slope (Bergoeing 2007). Cerro Chirripó is the highest summit in the country with an elevation of 3821 m (Alfaro and Gamboa 1999).

At approximately 3000 m elevation, the montane rainforest is replaced by the páramo ecosystem, which dominates the upper areas of the PNC. The páramo has an approximate extent of 6000 ha and temperatures can drop to −5 °C (Alfaro and Gamboa 1999). Páramo vegetation typically is composed of Dwarf Bamboo, *Chusquea subtessellata* Hitchcock, and there are small brooks, small, exposed boggy sites, and rock outcrops (Holz and Gradstein 2005). Marshes, meadows, expanses of bunch grass or bamboo, and thickets of shrubs, all are common in the páramo (Hooper and Carleton 1976).

We captured the mouse with a butterfly net and took measurements with calipers to the nearest 1 mm. We measured total head–body length, tail length, ear length, and foot length. We weighed the mouse with a spring scale. We compared eye sizes using photographs of the two species of *Scotinomys* occurring in Costa Rica. In addition, we reviewed records in the Global Biodiversity Information Facility (GBIF 2021), which mainly consisted of records from the Museum of Zoology of the University of Michigan (UMMZ), as well as the literature (Hooper 1972; Hooper and Carleton 1976; Reid 2009; Pasch et al. 2013; Naylor and Roach 2016; Bradley 2017).

To estimate the potential area occupied by *S. xerampelinus* in Costa Rica and Panama, we first calculated the area between 2100 and 3400 m elevation in the Cordillera de Talamanca and Irazú and Turrialba volcanoes

in the Cordillera Central. These upper and lower elevational limits were based on historical records in the literature for *S. xerampelinus*. However, published elevations are only presence data and may not represent the actual elevational limits of this species. Second, we estimated the area within an elevational range of 3400–3820 m, which considers the extension of the upper elevational range in one part of the extremely limited extent of occurrence of this species (Naylor and Roach 2016). To estimate the extent of these areas of distribution, both previously known as well as our hypothesized increase vertical range, we used the topographic layers for Costa Rica (Ortíz-Malavassi 2015) and Panama (http://www.diva-gis.org/gdata). For the Costa Rican analysis, we filtered by elevation data from a vector shapefile, from 2100 to 3400 m and higher than 3400 m, and the area of each subset then estimated. For the analysis of Panamanian geographic data, we transformed a raster.vrt file into a vector shapefile. We extracted the 2100 and 3400 m contour lines, made a polygon from these limits, and then calculated the polygon's area. Both Costa Rican and Panamanian data sets were examined at the same scale to avoid errors resulting from varying scales. We used the QGIS v. 3.10.6-A (Coruña) for this geographical analysis (QGIS 2020).

Results

***Scotinomys xerampelinus* (Bangs, 1902)**

Figure 1

New record. COSTA RICA – **Limón** • Cordillera de Talamanca, Parque Nacional Chirripó, Cerro Chirripó; 09°29′04″N, 083°29′19″W; 3820 m a.s.l.; 02.VIII.2015; Jairo García Céspedes leg; highest point of the Cerro; individual measured but not collected; 1 adult, sex undetermined.

Comments. JGC measured and photographed a *S. xerampelinus* found at rocky terrain dominated by Dwarf Bamboo (Fig. 2). The mouse was at an elevation of 3820 m, the highest point of altitude in Parque Nacional Chirripó (Fig. 3). The mouse prowled around the visitors maybe waiting for food. One tourist fed it a small piece of bread that the mouse started to eat immediately. In the

Figure 1. A Long-Tailed Singing Mouse *Scotinomys xerampelinus* at Cerro Chirripó at 3820 m elevation, Parque Nacional Chirripó, Costa Rica. Photo by Jairo García Céspedes.

Figure 2. Páramo habitat at Cerro Chirripó (3820 m elevation). The dominant grass looking plant is Dwarf Bamboo, *Chusquea subtessellata*. Parque Nacional Chirripó, Talamanca, Costa Rica. Photo by Jairo García Céspedes.

vicinity, some meters below the highest point of Cerro Chirripó, we observed two additional mice of the same species.

Other data records. Individuals of Long-tailed Singing Mouse have been collected or seen at numerous localities throughout its geographic range. Some extreme elevation records for this species found online at the UMMZ and other collections in the GBIF database (GBIF 2021) are from Cerro de la Muerte (Costa Rica), at 3100 m elevation (UMMZ 111773, 112258, and at least nine more records), and five localities in the Cordillera de Talamanca, including Villa Mills and Cerro Asunción (Costa Rica). There are also several localities by the Turrialba and Irazú volcanoes (Costa Rica) up to 3300 m a.s.l. (UMMZ 116836 and at least 57 more records). Eleven additional localities are from lower elevations. Pasch et al. (2013) hypothesized that the elevational range extends to 3451 m in Cerro de la Muerte, 3432 m in Volcán Irazú, and 3474 in Volcán Barú (Panama).

Identification. The mouse observed at Cerro Chirripó weighted 14 g, had a head–body length of 82 mm, a tail length of 68 mm, ear length of 16 mm, and foot length of 19 mm; these measurements agree with those of *S. xerampelinus* (Reid 2009; Villalobos-Chaves et al. 2016). This chocolate-brown mouse differs from the congeneric *S. teguina* by having a proportionally longer tail, which is about 80–85% of the head–body length; longer fur, which appears fuzzy rather than smooth; and smaller eyes (Reid 2009).

The potential area occupied by *Scotinomys xerampelinus* at elevations between 2100 and 3400 m is about 2101 km^2 in Costa Rica and 557 km^2 in Panamá, for a total of 2658 km^2. Including the new locality at 3820 m, the potential distribution area increases by 53 km^2 to 2711 km^2 (Fig. 3).

Discussion

Although *Scotinomys xerampelinus* is hypothesized to be common (Reid 2009), Naylor and Roach (2016) correctly pointed out that the extent of occurrence of this species is probably less than 10,000 km^2. Despite this restricted distribution, the species is Least Concern "in view of its presumed large and stable population, occurrence in a number of protected areas, and because it does not appear to be under threat and is unlikely to be declining at nearly the rate required to qualify for listing in a threatened category" (Naylor and Roach 2016). However, based on our elevational range analysis and on museum records, we hypothesize that the area occupied by *S. xerampelinus* could be as small as 2711 km^2 (Fig. 3). Outside of the protected areas, there are threats to this species, including habitat loss and the increase of mesocarnivores such as Coyote, *Canis latrans* (Say, 1823), Tayra *Eira barbara* (Linnaeus, 1758), and other predators. The increase in predators, which may negatively impact small mammals such as *S. xerampelinus*, is due to human-induced habitat changes (Mora and Ruedas 2018; Hody et al. 2019).



Figure 3. New altitudinal record (black dot) for the Long-tailed Singing Mouse (*Scotinomys xerampelinus*) at the top of Cerro Chirripó, Costa Rica. The yellow areas indicate lands between 2100 and 3400 m elevation, and the red areas indicate lands between 3400 and 3821 m elevation. Map by Gerardo Chaves.

Scotinomys xerampelinus previously was unknown from elevations above 3400 m (Reid 2009) and not reported from the top of Cerro Chirripó (Mora 1994; Carrillo et al. 2005; Chaverri 2008; Kappelle and Horn 2016). This could be due to a sampling bias or to an upward expansion of the species' range in recent decades. Because the fauna of Cerro Chirripó has been intensively studied (Kappelle and Horn 2016), sampling bias is an unlikely explanation for the previous lack of records from this site. The upslope shift is more likely the result, caused by increasing temperatures from climate change, which has forced several Costa Rican vertebrate species to move their distributions to higher elevations (Pounds et al. 1999). There is a general trend toward poleward and upward shifts of latitudinal and elevational boundaries of the ranges of several species (Chen et al. 2011; Rowe et al. 2015). Species inhabiting mountains may be more sensitive to climate change due to the various factors that affect community assemblages across elevational gradients (Neate-Clegg et al. 2018).

We point out the necessity for increasing systematic sampling for small mammals in the Cordillera de Talamanca, mainly at high elevations at Cerro Chirripó, Cerro Kamuk, Cerro Ventisqueros, Cerro Cuericí, Terbi, Los Crestones, Sabana de los leones, Valle de los Conejos and Fila Urán. Our new record for *S. xerampelinus* offers evidence that the new site, at 3820 elevation, currently is ecologically suitable for this species.

Acknowledgements

JMM and LIL acknowledge Daniel Tobias of the Unidad de Ciencias Básicas, Sede Atenas, and Emilce Rivera of Gestión Ecoturística, Sede Central, Universidad Técnica Nacional, for assigning time to work on this paper. We thank D. Villalobos-Chaves, an anonymous reviewer, and especially H.E. Ramirez-Chaves, Luis A. Ruedas and Robert Forsyth for their corrections and suggestions to improve this work.

Authors' Contributions

JMM identified the species and elaborated the manuscript. JGC found the specimen at the new locality and collected the data; LIL explored databases and helped on the manuscript preparation. GC conducted the

geographical analysis and elaborated the map. All four authors read, edited, and approved the manuscript.

References

Alfaro A (1897) Mamíferos de Costa Rica. Tipografía Nacional, San José, Costa Rica, 51 pp.

Alfaro E, Gamboa B (1999) Plantas comunes del Parque Nacional Chirripó. INBio, Santo Domingo de Heredia, Costa Rica, 283 pp.

Allen AP, Brown JH, Gilloley JF (2002) Global biodiversity, biochemical kinetics, and the Energetic-Equivalence Rule. Science 297: 1545–1548. https://doi.org/10.1126/science.1072380

Bergoeing JP (2007) Geomorfología de Costa Rica. Librería Francesa, San José, Costa Rica, 328 pp.

Bradley RD (2017) Long-Tailed Singing Mice mouse *Scotinomys xerampelinus*. In: Wilson DE, Lacher TE, Mittermeier RA (Eds.) Handbook of the mammals of the world. Volume 7. Rodents II. Lynx Editions, Barcelona, Spain, 366.

Brown JH (2001) Mammals on mountainsides: elevational patterns of diversity. Global Ecology and Biogeography 10: 101–109. https://doi.org/10.1046/j.1466-822x.2001.00228.x

Carrillo E, Wong G, Sáenz JC (2005) Mamíferos de los páramos de Costa Rica. In: Kappelle M, Horn SP (Eds.) Páramos de Costa Rica. INBio, Santo Domingo de Heredia, Costa Rica, 533–545.

Chaverri A (2008) Historia natural del Parque Nacional Chirripó, Costa Rica. INBio, Santo Domingo de Heredia, Costa Rica, 141 pp.

Chen I, Hill JK, Ohlemüller R, Roy DB, Thomas CD (2011) Rapid range shifts of species associated with high levels of climate warming. Science 333: 1024–1026. https://doi.org/10.1126/science.1206432

Dyer EE, Redding DW, Cassey P, Collen B, Blackburn TM (2020) Evidence for Rapoport's rule and latitudinal patterns in the global distribution and diversity of alien bird species. Journal of Biogeography 47 (6): 1362–1372. https://doi.org/10.1111/jbi.13825

Frantzius AV (1869) Los mamíferos de Costa Rica. Traducido por D. R. Cortéz. 1963. Revista de los Archivos Nacionales de Costa Rica 27: 237–281.

GBIF.org (2020) GBIF occurrence data. https://www.gbif.org/occurrence/search?q=Scotinomys%20xerampelinus. Accessed on: 2020-09-10.

Goodwin GG (1946) Mammals of Costa Rica. Bulletin of the American Museum of Natural History 87: 271–473.

Hillebrand H (2004) On the generality of the latitudinal diversity gradient. American Naturalist 163: 192–211. https://doi.org/10.1086/381004

Hody AW, Moreno R, Meyer NFV, Pacifici K, Kays R (2019) Canid collision—expanding populations of coyotes (*Canis latrans*) and crab-eating foxes (*Cerdocyon thous*) meet up in Panama. Journal of Mammalogy 11: 1819–1830. https://doi.org/10.1093/jmammal/gyz158

Holz I, Gradstein SR (2005) Phytogeography of the bryophyte floras of oak forests and páramo of the Cordillera de Talamanca, Costa Rica. Journal of Biogeography 32: 1591–1609. https://doi.org/10.1111/j.1365-2699.2005.01250.x

Hooper, ET (1972) A synopsis of the rodent genus *Scotinomys*. Occasional papers of the Museum of Zoology University of Michigan 665: 1–32.

Hooper ET, Carleton M (1976) Reproduction, growth and development in two contiguously allopatric rodent species, genus *Scotynomys*. Miscellaneous Publications Museum of Zoology University of Michigan 151: 1–52.

Kappelle M, Horn SP (2016) The páramo ecosystem of Costa Rica's highlands. In: Kappelle M (Ed.). Costa Rican ecosystems. University of Chicago Press, Chicago, USA, 492–523.

Kontopoulos D, Smith TP, Barraclough TG, Pawar S (2020) Adaptive evolution shapes the present-day distribution of the thermal sensitivity of population growth rate. PLoS Biology 18 (10): e3000894. https://doi.org/10.1371/journal.pbio.3000894

McCain CM (2007) Area and mammalian elevational diversity. Ecology 88 (1): 76–86. https://doi.org/10.1890/0012-9658(2007)88[76:aamed]2.0.co;2

McPherson AB (1971a) A new subspecies of squirrel *Sciurus variegatoides* Ogilby from Costa Rica. Revista de Biología Tropical 19 (1–2): 191–194.

McPherson AB (1971b) An analysis of factors influencing the distribution of Costa Rican Rodents. PhD thesis, Southern Illinois University, Carbondale, USA, 278 pp.

McPherson AB (1985) A biogeographical analysis of factors influencing the distribution of Costa Rican rodents. Brenesia 23: 97–273.

Mora JM (1994). Evaluación de fauna en el Parque Nacional Chirripó para la elaboración del plan General de Manejo. FUNDEVI-ICT, San José, Costa Rica, 11 pp.

Mora JM (2000) Los mamíferos silvestres de Costa Rica. Editorial UNED, San José, Costa Rica, 240 pp.

Mora JM, López LI (2014) Cambios en la distribución de las aves: ¿Qué puerta hemos abierto?; los casos del chicopiojo (*Campylurhynchus rufinucha*) y la paloma ala blanca (*Zenaida asiatica*) en Costa Rica. Ceiba 52 (2): 230–236. https://doi.org/10.5377/ceiba.v52i2.1760

Mora JM, Ruedas LA (2018) *Sylvilagus dicei* Harris, 1932. In: Smith AT, Johnston CH, Alves PC, Hackländer K (Eds.) Lagomorphs: pikas, rabbits, and hares of the world. The Johns Hopkins University Press, Baltimore, USA, 135–137.

Mora JM, López LI, Sánchez R, Ruedas LA (2020) Murciélagos de la Reserva Biológica Alberto Manuel Brenes. Coordinación de Investigación, Sede de Occidente, Universidad de Costa Rica, San Ramón, Costa Ria, 179 pp.

Musser GG, Carleton MD (2005) Superfamily Muroidea. In: Wilson DE, Reeder DA (Eds.) Mammal species of the world: a geographic and taxonomic reference. The John Hopkins University Press, Baltimore, USA, 894–1531.

Naylor L, Roach N (2016) *Scotinomys xerampelinus*. The IUCN Red List of threatened species. https://doi.org/10.2305/IUCN.UK.2016-2.RLTS.T20053A22389972.en. Accessed on: 2020-6-14.

Neate-Clegg MHC, Jones SEI, Burdekin O, Jocque M, Sekercioglu CH (2018) Elevational changes in the avian community of a Mesoamerican cloud forest park. Biotropica 50 (5): 805–815. https://doi.org/10.1111/btp.12596

Ortíz-Malavassi E (2015) Atlas digital de Costa Rica. Investiga TEC 23: 24.

Pasch B, Bolker BM, Phelps SM (2013) Interspecific dominance via vocal interactions mediates altitudinal zonation in Neotropical singing mice. The American Naturalist 182: E161–E173. https://doi.org/10.1086/673263

Pianka ER (1989) Latitudinal gradients in species diversity. Trends in Ecology & Evolution 4: 223. https://doi.org/10.1016/0169-5347(89)90163-8

Pounds JA, Fogden MPL, Campbell JH (1999) Biological response to climate change on a tropical mountain. Nature 398 (6728): 611–615. https://doi.org/10.1038/19297

QGIS Development Team (2020) QGIS Geographic Information System. http://qgis.osgeo.org. Accessed on: 2020-6-10.

Ramírez-Fernández JD, Durán FJ, Fernández-Vargas M (2020) First record of Tweedy's Crab-eating Rat, *Ichthyomys tweedii* Anthony, 1921 (Rodentia, Cricetidae, Sigmodontinae), for Costa Rica. Check List 16 (2): 435–440. https://doi.org/10.15560/16.2.435

Reid F (2009) A field guide to the mammals of Central America and southeast Mexico, 2nd edition. Oxford University Press, New York, USA, 334 pp.

Rodríguez-Herrera B, Wilson DE, Fernández M, Pineda W (2005) La mastozoología en Costa Rica: historia, recolecta, localidades y composición de especies. Brenesia 63-64: 89–112.

Rodríguez-Herrera B, Chinchilla FA, May-Collado L (2002) Lista de

especies, endemismo y conservación de los mamíferos de Costa Rica. Revista Mexicana de Mastozoología 6 (1): 21–57.

Rodríguez-Herrera, B, Ramírez-Fernández JD, Villalobos-Chaves D, Sánchez R (2014) Actualización de la lista de especies de mamíferos vivientes de Costa Rica. Mastozoología Neotropical 21 (2): 275–289.

Rowe KC, Rowe KMC, Tingley MW, Koo MS, Patton JL, Conroy G, Perrine JD, Beissinger SR, Moritz C (2015) Spatially heterogeneous impact of climate change on small mammals of montane California. Proceedings of the Royal Society B 282: 20141857. https://doi.org/10.1098/rspb.2014.1857

Ryan RM (1963) The biotic provinces of Central America as indicated by mammalian distribution. Acta Zoológica Mexicana 6: 1–54.

SINAC Sistema Nacional de Áreas de Conservación (2014) V Informe Nacional al Convenio sobre la Diversidad Biológica, Costa Rica. GEF-PNUD, San José, Costa Rica, 192 pp.

Villalobos-Chaves D, Ramírez-Fernández JD, Chacón-Madrigal E, Pineda-Lizano W, Rodríguez-Herrera B (2016) Clave para la identificación de los roedores de Costa Rica. Universidad de Costa Rica, Escuela de Biología, San José, Costa Rica, 37 pp.

Veas-Ayala N, Quesada-Román A, Hidalgo HG, Alfaro EJ (2018) Humedales del Parque Nacional Chirripó, Costa Rica: características, relaciones geomorfológicas y escenarios de cambio climático. Revista de Biología Tropical 66 (4): 1436–1448. https://doi.org/10.15517/rbt.v66i4.31477

Wen Z, Feijó A, Cheng J, Du Y, Ge D, Xia L, Yang Q (2021) Explaining mammalian abundance and elevational range size with body mass and niche characteristics. Journal of Mammalogy 102: gyaa093. https://doi.org/10.1093/jmammal/gyaa093

Wilson DE (1983) Lista de Mamíferos de Costa Rica. In: Janzen DH (Ed.) Historia natural de Costa Rica. Editorial de la Universidad de Costa Rica, San José, Costa Rica, 457–514.

Wilson DE, Timm RM, Chinchilla FA (2002) Mamíferos de Costa Rica. In: Ceballos G, Simonetti JA (Eds.) Diversidad y conservación de los mamíferos neotropicales. CONABIO, México, 227–254.